

Case reports

Aortic atresia with double inlet left ventricle, rudimentary left sided right ventricle, and ventriculoarterial discordance

CHARLES A BULLABOY, HERBERT S HARNED JR

From the Department of Pediatrics, Division of Pediatric Cardiology, Children's Hospital of the King's Daughters, Eastern Virginia Medical School, Norfolk, Virginia; and the Department of Pediatrics, Division of Pediatric Cardiology, University of North Carolina, School of Medicine, Chapel Hill, North Carolina, USA

SUMMARY Aortic atresia associated with ventriculoarterial discordance (transposition) was found at necropsy in a 3 month old neonate. This is a rare association.

Aortic atresia is one of the most common causes of congestive heart failure and death in early life.¹ Usually, the great arteries are normally related. We report the unusual association of aortic atresia with ventriculoarterial discordance.

Case report

A 3 day old male neonate was referred with a history of tachypnoea and cyanosis during feeding. The blood pressure on admission was 75/40 mm Hg in the right arm and 90/50 mm Hg in the right leg. Physical examination detected a right ventricular impulse, single second heart sound, a grade 2/6 systolic ejection murmur along the left sternal border, an S₃ gallop, and hepatomegaly. A chest radiograph showed a cardiothoracic ratio of 0.65 with increased pulmonary vascularity. The electrocardiogram showed right atrial and left ventricular hypertrophy. Arterial oxygen pressure was 51 mm Hg (6.8 kPa) in room air.

At cardiac catheterisation oximetry showed an increment in oxygen saturation at the right sided ventricular level with similar saturations in the pulmonary arteries and aorta. Pulmonary arterial pressure was equal to the systemic pressure. Ventriculography showed a large morphological left ventricle with ventriculoarterial discordance (Figure *ai* and *bi*). The

catheter entered the aorta from the pulmonary artery via a duct. Aortography showed a left sided anteriorly positioned hypoplastic ascending aorta (Figure *aii* and *bii*).

The patient died in 1974 at 3 months of age of refractory congestive heart failure.

At necropsy viscerotransposition was present. Systemic venous drainage was into the morphological right atrium. The morphological left atrium received the four pulmonary veins. The normally formed atrial septum had a patent foramen ovale. The right sided atrioventricular valve had two leaflets and the left sided atrioventricular valve was tricuspid; both entered the left ventricle (a double inlet left ventricle). The origin of the posterior pulmonary artery was from this chamber. A rudimentary left sided right ventricle gave the origin to the anterior and leftward atretic aortic valve and hypoplastic ascending aorta. The great arteries were joined by a duct.

Discussion

Aortic atresia and ventriculoarterial discordance (transposition) are rarely found in tandem. To our knowledge only five such cases have been reported. Two were associated with double inlet left ventricles of the left ventricular type,^{2,3} one with corrected transposition,⁴ one with complete transposition,⁵ and one with absent tricuspid valve with atrioventricular discordance.⁶

Systemic blood flow is rarely obstructed in hearts with ventriculoarterial discordance. A restrictive bul-

Requests for reprints to Dr Charles A Bullaboy, Division of Pediatric Cardiology, Children's Hospital of the King's Daughters, 800 W Olney Road, Norfolk, Virginia 23507, USA.




Figure (a) Lateral and (b) anteroposterior angiograms (i) from the ventricular injection showing filling of the pulmonary artery, which is rightward and posterior in position (no contrast exits into the aorta from this injection), and (ii) from the retrograde aortic injection showing filling of the pulmonary artery which is posterior to the hypoplastic ascending aorta; the aorta is leftward and anterior in relation to the pulmonary artery.

boventricular foramen and accessory endocardial cushion tissue are among the pathological mechanisms that limit aortic flow by producing subaortic stenosis or atresia.⁷ Why this occurs is a matter of speculation.

We thank Dr Glenn Rosenquist for examining the necropsy specimen.

References

- 1 Roberts WC, Perry LW, Chandra RS, Myers GE, Shapiro SR, Scott LP. Aortic valve atresia: a new classification based on necropsy study of 73 cases. *Am J Cardiol* 1976; 37: 753-6.
- 2 Van Praagh R, Pletl JA, Van Praagh S. Single ventricle. Pathology, embryology, terminology, and classification. *Herz* 1979; 4: 113-50.
- 3 Macartney FJ, Anderson RH. Angiocardiography and hemodynamics of the univentricular heart with two atrioventricular valves or a common atrioventricular valve. In: Anderson RH, Shinebourne EA, eds. *Pediatric cardiology* 1977. Edinburgh and London: Churchill Livingstone, 1978: 345-59.
- 4 Deanfield JE, Anderson RH, Macartney FJ. Aortic atresia with "corrected transposition of the great arteries" (atrioventricular and ventriculoarterial discordance). *Br Heart J* 1981; 46: 683-6.
- 5 McGarry KM, Taylor JFN, Macartney FJ. Aortic atresia occurring with complete transposition of great arteries. *Br Heart J* 1980; 44: 711-3.
- 6 Brenner JI, Bharati S, Winn WC Jr, Lev M. Absent tricuspid valve with aortic atresia in mixed levocardia (atria situs solitus, L-loop). A hitherto undescribed entity. *Circulation* 1978; 57: 836-40.
- 7 Freedom RM, Dische MR, Rowe RD. Pathologic anatomy of subaortic stenosis and atresia in the first year of life. *Am J Cardiol* 1977; 39: 1035-44.